



TITLE:

COMMUTING CONTRACTIONS の SIMULTANEOUS UNITARY DILATION(Linear Operators and Inequalities)

AUTHOR(S):

OKAYASU, Takateru

CITATION:

OKAYASU, Takateru. COMMUTING CONTRACTIONS の SIMULTANEOUS UNITARY
DILATION(Linear Operators and Inequalities). 数理解析研究所講究録 1994, 860: 7-11

ISSUE DATE:

1994-03

URL:

<http://hdl.handle.net/2433/83829>

RIGHT:

COMMUTING CONTRACTIONS の SIMULTANEOUS UNITARY DILATION

山形大学理学部 岡 安隆 照 (Takateru OKAYASU)

The following matter is really fundamental:

Sz.-Nagy's Unitary Dilation Theorem. Let T be a contraction on a Hilbert space \mathcal{H} . Then, there exist an enlarged Hilbert space $\mathcal{K} \supseteq \mathcal{H}$ and a unitary U , called a unitary dilation of T , on \mathcal{K} , such that

$$T^m = PU^m|_{\mathcal{H}} \quad \text{for } m = 0, 1, 2, \dots,$$

where P is the projection on \mathcal{K} onto \mathcal{H} .

This yields, and, is yielded by, the so-called

von Neumann Inequality. Let T be a contraction on a Hilbert space. Then,

$$\|p(T)\| \leq \|p\| = \sup_{z \in \mathbb{T}} |p(z)|$$

holds for any polynomial p with complex coefficients.

The “logical equivalence” is accompanied by the the following

Theorem [6]. If a set of commuting contractions on a Hilbert space \mathcal{H} , T_1, T_2, \dots, T_n , admits a simultaneous unitary dilation, namely, there exist a Hilbert space $\mathcal{K} \supseteq \mathcal{H}$ and commuting unitaries U_1, U_2, \dots, U_n on \mathcal{K} , such that

$$T_1^{m_1} T_2^{m_2} \dots T_n^{m_n} = P U_1^{m_1} U_2^{m_2} \dots U_n^{m_n} |_{\mathcal{H}}$$

for $m_1, m_2, \dots, m_n = 0, 1, 2, \dots$, where P is the projection on \mathcal{K} onto \mathcal{H} , then T_1, T_2, \dots, T_n enjoys the von Neumann inequality, namely,

$$\|(p_{ij}(T_1, T_2, \dots, T_n))\| \leq \|(p_{ij})\| = \sup_{z_1, z_2, \dots, z_n \in \mathbb{T}} \|(p_{ij}(z_1, z_2, \dots, z_n))\|$$

holds for any $m \times m$ matrix (p_{ij}) whose entries are polynomials with complex coefficients; and *vice versa*.

On the other hand, the following theorems are known:

Andô's Theorem [1]. Any pair of commuting contractions on a Hilbert space admits a simultaneous unitary dilation.

Andô's Theorem [2]. Any triple of commuting contractions on a Hilbert space, one of which double commutes with others, admits a simultaneous unitary dilation.

We, aside, have examples of triples of commuting contractions which do not admit a simultaneous unitary dilation, [4], [8] and [9].

In [6] we gave the following theorem and corollary:

Theorem. Suppose each of sets of commuting contractions, S_1, S_2, \dots, S_m and T_1, T_2, \dots, T_n , on a Hilbert space, admits a simultaneous unitary dilation, and every S_j double commutes with all T_k . If the set S_1, S_2, \dots, S_m generates a nuclear C^* algebra, then the set $S_1, S_2, \dots, S_m, T_1, T_2, \dots, T_n$ admits a simultaneous unitary dilation.

Collorary. Suppose S is a *GCR* contraction, i.e., a contraction which generates a *GCR* (postliminal) algebra, T_1, T_2, \dots, T_n commuting contractions, on a Hilbert space, the set T_1, T_2, \dots, T_n admits a simultaneous unitary

dilation. and S double commutes with all T_k . Then the set S, T_1, T_2, \dots, T_n admits a simultaneous unitary dilation.

The following, furthermore, turned out to be true [7]:

Theorem. Suppose each of sets of commuting contractions, S_1, S_2, \dots, S_m and T_1, T_2, \dots, T_n , on a Hilbert space, admits a simultaneous unitary dilation, and every S_j double commutes with all T_k . If the set S_1, S_2, \dots, S_m generates an injective von Neumann algebra, then the set $S_1, S_2, \dots, S_m, T_1, T_2, \dots, T_n$ admits a simultaneous unitary dilation.

Collorary. Suppose S is a type I contraction, i.e., a contraction which generates a type I von Neumann algebra, T_1, T_2, \dots, T_n commuting contractions, on a Hilbert space, the set T_1, T_2, \dots, T_n admits a simultaneous unitary dilation and S double commutes with all T_k . Then, the set S, T_1, T_2, \dots, T_n admits a simultaneous unitary dilation.

We here will improve the theorem, by making the assumption thin as the following

Theorem. Suppose each of sets of commuting contractions, S_1, S_2, \dots, S_m and T_1, T_2, \dots, T_n , on a Hilbert space, admits a simultaneous unitary dilation, and every S_j double commutes with all T_k . Then, the set $S_1, S_2, \dots, S_m, T_1, T_2, \dots, T_n$ admits a simultaneous unitary dilation.

This is the aimed theorem of ours. A proof of this is given, on account of the Steinspring representation of completely positive maps, by the preceding theorem and the

Arveson Theorem [3, Theorem 1.3.1]. Let \mathcal{H}, \mathcal{K} be Hilbert spaces, V a bounded operator from \mathcal{H} into \mathcal{K} , and \mathcal{B} a $*$ -subalgebra of $\mathcal{B}(\mathcal{K})$, the full operator algebra, which satisfies that $[\mathcal{B}V\mathcal{H}] = \mathcal{K}$. Then, for every $T \in (V^*\mathcal{B}V)'$ there exists a unique $\tilde{T} \in \mathcal{B}'$ such that $\tilde{T}V = VT$, and the mapping $(\) \mapsto (V^*\mathcal{B}V)'$ is a σ weakly continuous $*$ -homomorphism.

We have as well

Collorary. Suppose each of pairs of commuting contractions, S_1, S_2 , and T_1, T_2 , on a Hilbert space, admits a simultaneous unitary dilation, and each of S_1, S_2 double commutes with T_1, T_2 . Then, the set S_1, S_2, T_1, T_2 admits a simultaneous unitary dilation.

Our theorem, of course, gives a good understanding to Andô's "triple" assertion; on the Andô's "pair" assertion, the next matter sheds light:

Theorem [5, Theorem 6]. Let T be a contraction on a Hilbert space \mathcal{H} , U the minimal unitary dilation of T . Then for every $S \in \{T\}'$ there exists $\tilde{S} \in \{U\}'$ such that $S = P\tilde{S}|_{\mathcal{H}}$ and $\|\tilde{S}\| = \|S\|$.

References

1. T. Andô, On a pair of commuting contractions, *Acta Sci. Math.* 24(1963), 88-90.
2. T. Andô, Unitary dilation for a triple of commuting contractions, *Bull. Acad. Polonaise Math.* 24(1976), 851-853.
3. W. B. Arveson, Subalgebras of C^* algebras, *Acta Math.* 123(1969), 141-224.
4. M. J. Crabb, and A. M. Davie, von Neumann's inequality for Hilbert space operators, *Bull. London Math. Soc.* 7(1975), 49-50.
5. R. G. Douglas, P. S. Muhley, and C. Pearcy, Lifting commuting operators, *Michigan Math. Journ.* 15(1968), 385-395.
6. T. Okayasu, The von Neumann inequality and dilation theorems for contractions, *Operator Theory: Adv. Appl.* 59(1992), 285-291.
7. 岡安隆照, The von Neumann inequality, 日本数学会秋期総合分科会, 1992/10/6.

8. I. Parrott, Unitary dilations for commuting contractions, *Pacif. Journ. Math.* 34(1970), 481-490.
9. N. Th. Varopoulos, On an inequality of von Neumann and an application of metric theory of tensor products to operator theory, *Journ. Funct. Anal.* 16(1974), 83-100.